

Fascinating Life Sciences

Valentí Rull
Ana Carolina Carnaval
Editors

Neotropical Diversification: Patterns and Processes



Springer

Fascinating Life Sciences

This interdisciplinary series brings together the most essential and captivating topics in the life sciences. They range from the plant sciences to zoology, from the microbiome to macrobiome, and from basic biology to biotechnology. The series not only highlights fascinating research; it also discusses major challenges associated with the life sciences and related disciplines and outlines future research directions. Individual volumes provide in-depth information, are richly illustrated with photographs, illustrations, and maps, and feature suggestions for further reading or glossaries where appropriate.

Interested researchers in all areas of the life sciences, as well as biology enthusiasts, will find the series' interdisciplinary focus and highly readable volumes especially appealing.

More information about this series at <http://www.springer.com/series/15408>

Valentí Rull • Ana Carolina Carnaval
Editors

Neotropical Diversification: Patterns and Processes

 Springer

Editors

Valentí Rull
Laboratory of Paleoecology
Institute of Earth Sciences Jaume Almera
(CSIC)
Barcelona, Spain

Ana Carolina Carnaval
Department of Biology
City College of New York
New York, NY, USA

Biology Program, Graduate Center of CUNY
New York, NY, USA

ISSN 2509-6745

Fascinating Life Sciences

ISBN 978-3-030-31166-7

<https://doi.org/10.1007/978-3-030-31167-4>

ISSN 2509-6753 (electronic)

ISBN 978-3-030-31167-4 (eBook)

© Springer Nature Switzerland AG 2020

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG.
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Acknowledgements

The editors are very grateful to all the authors who contributed to this book for their dedication and hard work leading to high-quality chapters. We also wish to thank the reviewers for their time and expertise (in alphabetic order): Alexandre Antonelli, Chrisitine Bacon, John M. Bates, João M. Capurucho, Rosane Collevatti, Guarino Colli, Chris Dick, José A. Diniz Filho, Leandro Duarte, Joost F. Duivervoorden, José M. Fernández Palacios, Stephen Hopper, Marie-Pierre Ledru, Peter Linder, Francis Mayle, Fabian Michelangeli, Cristina Miyaki, Toby Pennington, Ivan Prates, Renato Recoder, Marcelo Reginato, Camila Ribas, Ignacio de la Riva, Miguel T. Rodrigues, Fernando Silveira, Beryl Simpson and Ángel Viloría. We greatly acknowledge the invitation of Lars Koerner, from Springer, for editing/writing the book and Sivachandran Ramanan for his constant support in editorial and production tasks.

Contents

1	Introduction	1
	Valentí Rull and Ana Carolina Carnaval	
Part I Neotropical Diversity and Diversification		
2	Neotropical Diversification: Historical Overview and Conceptual Insights	13
	Valentí Rull	
3	Beyond Refugia: New Insights on Quaternary Climate Variation and the Evolution of Biotic Diversity in Tropical South America	51
	Paul A. Baker, Sherilyn C. Fritz, David S. Battisti, Christopher W. Dick, Oscar M. Vargas, Gregory P. Asner, Roberta E. Martin, Alexander Wheatley, and Ivan Prates	
4	Tropical and Subtropical South America: A Study of Community Turnover Across Environmental Gradients	71
	João Claudio Sousa Nascimento, Millke Jasmine Arminini Morales, Wendy Yohana Arroyo-Pérez, Juliana José, Monica Paiva Quast, and Vera Nisaka Solferini	
5	Evolutionary Macroecology and the Geographical Patterns of Neotropical Diversification	85
	Fabricio Villalobos, Jesús N. Pinto-Ledezma, and José Alexandre Felizola Diniz-Filho	
6	Evolutionary Imprints on Species Distribution Patterns Across the Neotropics	103
	Renan Maestri and Leandro Duarte	
7	Diatom Diversity and Biogeography Across Tropical South America	121
	Xavier Benito and Sherilyn C. Fritz	

8	Avian Diversity in Humid Tropical and Subtropical South American Forests, with a Discussion About Their Related Climatic and Geological Underpinnings	145
	Manuel Nores	
9	Contrasting Patterns of Temporal Diversification in Neotropical Butterflies: An Overview	189
	Andrew V. Z. Brower and Ivonne J. Garzón-Orduña	
Part II Regional Biodiversity Patterns and Diversification Processes		
10	The Origin and Evolution of Amazonian Species Diversity	225
	J. Cracraft, C. Camila Ribas, F. M. d’Horta, J. Bates, R. P. Almeida, A. Aleixo, J. P. Boubli, K. E. Campbell, F. W. Cruz, M. Ferreira, S. C. Fritz, C. H. Grohmann, E. M. Latrubesse, L. G. Lohmann, L. J. Musher, A. Nogueira, A. O. Sawakuchi, and P. Baker	
11	Patterns and Processes of Diversification in Amazonian White Sand Ecosystems: Insights from Birds and Plants	245
	J. M. G. Capurucho, S. H. Borges, C. Cornelius, A. Vicentini, E. M. B. Prata, F. M. Costa, P. Campos, A. O. Sawakuchi, F. Rodrigues, A. Zular, A. Aleixo, J. M. Bates, and C. Camila Ribas	
12	On the Young Savannas in the Land of Ancient Forests	271
	Josué A. R. Azevedo, Rosane G. Collevatti, Carlos A. Jaramillo, Caroline A. E. Strömberg, Thaís B. Guedes, Pável Matos-Maraví, Christine D. Bacon, Juan David Carrillo, Søren Faurby, and Alexandre Antonelli	
13	Diversity, Endemism, and Evolutionary History of Montane Biotas Outside the Andean Region	299
	Thaís B. Guedes, Josué A. R. Azevedo, Christine D. Bacon, Diogo B. Provete, and Alexandre Antonelli	
14	Diversification in Ancient and Nutrient-Poor Neotropical Ecosystems: How Geological and Climatic Buffering Shaped Plant Diversity in Some of the World’s Neglected Hotspots	329
	Fernando A. O. Silveira, Roberta L. C. Dayrell, Cecilia F. Fiorini, Daniel Negreiros, and Eduardo L. Borba	
15	The Pantepui “Lost World”: Towards a Biogeographical, Ecological and Evolutionary Synthesis of a Pristine Neotropical Sky-Island Archipelago	369
	Valentí Rull and Teresa Vegas-Vilarrúbia	

16	Patterns of Species and Lineage Diversity in the Atlantic Rainforest of Brazil	415
	Elen A. Peres, Ricardo Pinto-da-Rocha, Lúcia G. Lohmann, Fabian A. Michelangeli, Cristina Y. Miyaki, and Ana Carolina Carnaval	
17	Forests Diversity in the Mexican Neotropics: A Paleoecological View	449
	Blanca L. Figueroa-Rangel, Miguel Olvera-Vargas, Socorro Lozano-García, Gerald Islebe, Nuria Torrescano, Susana Sosa-Najera, and Ana P. Del Castillo-Batista	
18	Bioregions of Eastern Brazil, Based on Vascular Plant Occurrence Data	475
	Marcelo Reginato and Fabián A. Michelangeli	
19	Human Contribution to Amazonian Plant Diversity: Legacy of Pre-Columbian Land Use in Modern Plant Communities	495
	Encarni Montoya, Umberto Lombardo, Carolina Levis, Gerardo A. Aymard, and Francis E. Mayle	
20	Historical Biogeography of Caribbean Plants Revises Regional Paleogeography	521
	Julissa Roncal, María Esther Nieto-Blázquez, Agustín Cardona, and Christine D. Bacon	
21	The Diversification of Extant Angiosperms in the South America Dry Diagonal	547
	Rosane G. Collevatti, Natácia E. Lima, and Luciana C. Vitorino	
Part III Taxon-Based Local and Regional Studies		
22	Amphibians and Reptiles of Venezuelan Guayana: Diversity, Biogeography and Conservation	571
	Celsa Señaris and Fernando J. M. Rojas-Runjaic	
23	Origin of Elevational Replacements in a Clade of Nearly Flightless Birds: Most Diversity in Tropical Mountains Accumulates via Secondary Contact Following Allopatric Speciation	635
	Carlos Daniel Cadena and Laura N. Céspedes	
24	Clade-Specific Biogeographic History and Climatic Niche Shifts of the Southern Andean-Southern Brazilian Disjunction in Plants	661
	Federico Luebert, Maximilian Lörch, Rafael Acuña, Renato Mello-Silva, Maximilian Weigend, and Jens Mutke	

25	Biotic and Landscape Evolution in an Amazonian Contact Zone: Insights from the Herpetofauna of the Tapajós River Basin, Brazil	683
	Leandro J. C. L. Moraes, Camila C. Ribas, Dante Pavan, and Fernanda P. Werneck	
26	Diversification Processes in Lizards and Snakes from the Middle São Francisco River Dune Region, Brazil	713
	Renato Sousa Recoder and Miguel Trefaut Rodrigues	
27	Unexpected Beta-Diversity Radiations in Highland Clades of Andean Terraranae Frogs	741
	Ignacio De la Riva	
28	Disparate Patterns of Diversification Within Liolaemini Lizards . . .	765
	Melisa Olave, Andrea González Marín, Luciano J. Avila, Jack W. Sites Jr., and Mariana Morando	
29	Diversification History of Neotropical Lecythidaceae, an Ecologically Dominant Tree Family of Amazon Rain Forest	791
	Oscar M. Vargas and Christopher W. Dick	
Part IV Epilogue		
30	Conservation in the Neotropics: A Final Reflection	813
	Ana Carolina Carnaval	

Chapter 1

Introduction



Valentí Rull and Ana Carolina Carnaval

This book addresses the colossal biodiversity of the Neotropics and provides a range of potential explanations for its origin and maintenance. In general, we use the term “Neotropics” as synonym of “tropical America”—i.e., the part of this continent situated between the tropics of Cancer and Capricorn—rather than of the Neotropical biogeographical realm, which includes also the southern part of South America and excludes the Mexican highlands. However, the geographical distribution of the taxonomic groups studied in some chapters makes necessary to consider certain extratropical areas for a thorough study.

The main differential feature of this book with respect to other compilations and reviews on the same subject is that it is not attached to any particular idea on how the Neotropical biodiversity has originated. Instead, it considers that there is no a single explanation applicable to any Neotropical region or taxonomic group. We (the editors) are convinced that different environmental drivers acting at multiple spatio-temporal scales have led to the complex environmental-evolutionary-ecological interactions that resulted in current biogeographical patterns (Rull 2011), and this compilation is a testimony of that.

For this reason, we have invited scholars who are working on the subject regardless their personal views on Neotropical diversification. Each author has been free to choose the topic of his/her paper and to develop it according to their own preferences, without any formal or conceptual constraint besides the editorial author’s instructions. As a result, the book gathers as many views as possible on the

V. Rull (✉)

Laboratory of Paleocology, Institute of Earth Sciences Jaume Almera (ICTJA-CSIC),
Barcelona, Spain
e-mail: vrull@ictja.csic.es

A. C. Carnaval

Department of Biology, City College of New York, New York, NY, USA

Biology Program, Graduate Center of CUNY, New York, NY, USA

© Springer Nature Switzerland AG 2020

V. Rull, A. C. Carnaval (eds.), *Neotropical Diversification: Patterns and Processes*,
Fascinating Life Sciences, https://doi.org/10.1007/978-3-030-31167-4_1

1

causes of the extant Neotropical biodiversity, according to the awareness of the editors and the availability of the contributors.

It could be argued that this could create confusion, rather than help the reader to acquire a sound view of the status of the problem. We do not agree with that opinion, for several reasons. After more than three centuries of study, it is clear that the origin of Neotropical biodiversity and of large-scale diversity patterns (e.g., the latitudinal diversity gradients) are not simple problems with a single explanation (Brown 2014). Any attempt to reduce global macroecological patterns to a particular cause runs the risk of being inaccurate or unrealistic. It may be that a general theory on Neotropical biodiversity is just a human expectation or construct, and that explanations should be explored case by case, biome by biome, or region by region (Carnaval et al. 2014). Syntheses such as the presented here may be a good starting point to test these ideas. Avoiding the potential pitfalls of focusing the narrative on one or few sensationalist messages, regardless the soundness of its empirical support—a not unusual practice in current scientific literature—we created space for multiple views and data, allowing the contributing scholars to build and share their own perspective on Neotropical diversity and diversification. We consider this to be more useful for those readers interested in figuring out the true state of the art.

1 Book Plan

The book has been subdivided into three main parts: (I) Neotropical diversity and diversification, (II) regional biodiversity patterns and diversification processes, (III) taxon-based studies at the local and regional scales. An epilogue links the topics discussed here with the conservation of Neotropical biodiversity. The first part examines diversity patterns and diversification processes at a Neotropical scale using multitaxon approaches or broad taxonomic categories, such as plants, diatoms or birds, among others. The second part gathers chapters focused on more specific but large-scale Neotropical regions. Diversity patterns and diversification processes have been most intensely studied in the Amazon, and this region is often used as the preferred model for discussing the origin and maintenance of Neotropical biodiversity. In Part II, however, we ensured to include areas that go beyond the Amazon basin, considering, for example, the Atlantic rainforests, the Mexican forests, and the Caribbean, a region that is rarely contemplated in studies about Neotropical diversification. The third section of the book includes studies of particular taxonomic groups (amphibians, reptiles, flightless birds, angiosperm families) on specific Neotropical areas such as the Guayanan and the Andean highlands, or selected river catchments. Below, we briefly summarize the content of chapters of the main book sections.

Part I: Neotropical Diversity and Diversification

This part is composed of eight chapters.

Rull introduces the focus area of this book, and discusses Neotropical diversification from historical and conceptual perspectives. Four historical steps are recognized, namely the discovery of the latitudinal biodiversity gradients, the

biogeographic phase, the paleoecological phase, and the phylogeographical phase. Conceptually, this chapter explores the main handicaps that have hindered the progress towards an explanation for the comparatively high Neotropical biodiversity, with emphasis on the shifting from one paradigm to another, the geographical and taxonomic extrapolation of particular case studies, the selection of biased evidence to support different hypotheses, the identification of Pleistocene diversification with the refugial hypothesis, and the inference of diversification mechanisms from solely diversification timing. Some solutions are proposed to circumvent these drawbacks.

Baker et al. follow Rull's opening by revisiting the Haffer's refuge hypothesis in light of the paleoclimatic evidence gathered during the last 50 years. Their message to us is that the role of Quaternary climatic changes on Neotropical diversification should be reevaluated. The authors emphasize two main features that should be considered in addition to the glacial-interglacial cyclicality, namely the existence of an east-west precipitation dipole due to precessional cycles of ~20-kyr period, and the occurrence of phases of anomalously high rainfall of ~1500-y recurrence consistent with cold North-Atlantic stadials (Dansgaard-Oeschger cycles). The combination of these climatic modes conform a much more complex scenario of climatic change in time and space than originally proposed by the refuge hypothesis.

The biological significance of climatic transitions, this time in space, is the focus of the next chapter. Nascimento et al. investigate the effect of the Tropic of Capricorn in the distribution of Neotropical communities, particularly comparing the changes observed in the composition of biological communities in geographical space. By cleaning and analyzing nearly 15 million records of 124,066 species, all available online, they show that only in the cis-Andean region one can observe a latitudinal gradient in the degree of species turnover. Contrary to previous taxon-specific analyses, this chapter suggests that the tropical-subtropical transition does not influence the rate at which turnover indexes change in South America—at least not when data from multiple groups are combined. Although the data indicate that the replacement of species is largely responsible for changes in community composition, the Tropic of Capricorn does not seem to represent a common marker of concerted changes in community turnover at this large scale.

Villalobos et al. also focus on large-scale patterns of biodiversity in geographic space, but they utilize new methods in the field of macroecology to incorporate evolutionary thinking into studies of species ranges. They illustrate these novel approaches with data from the Furnariidae, a model group of Neotropical birds. Their contribution shows that multiple metrics now enable the inclusion of speciation and extinction processes into data analyses and provide coherent results and insights. The chapter ends with a discussing of ways to expand the field by incorporating lineage dispersal over evolutionary time as a major contributor to geographic diversity gradients, and the promise of simulation studies in historical inference.

Similar to Nascimento et al., Maestri and collaborators combine data from thousands of species and explore possible correlates of biodiversity patterns at a continental scale. Yet, they specifically explore how the spatial distribution of

evolutionary history compare across major groups of organisms. For that, they combine phylogenetic data and occurrence information from 1100 + species of vertebrates whose evolutionary histories are primarily confined to the Neotropics. They find that the spatial patterns of phylogenetic dissimilarity between areas are very similar across groups, despite the fact that they have different ecological requirements and ages. Their analysis also suggests that biogeographical relationships across regions explain phylogenetic similarity patterns better than present-day environments.

Nores discusses the biogeographical patterns observed in the birds of tropical and subtropical South America. Using range data, he introduces and discusses major geological and climatic events in the Neogene and the Pleistocene that are believed to have impacted these fauna, including the uplift of the Andes (splitting pan-Amazonia into a trans- and a cis-Andean regions, and leading to the formation of large Amazonian rivers), and the closing of the Isthmus of Panama (allowing for faunal interchange, increase in diversity, and new radiations). He argues that Amazonia and other tropical forests remained predominantly forested over time, even during glacial periods, and claims that marine transgressions in the Tertiary and lower Quaternary, tied to Amazonian rivers, were important by fragmenting species ranges and leading to speciation. Nores ends this chapter alerting us to the significant drop in species diversity and abundance recently observed in South America, particularly in protected areas of Amazonia, the lower Eastern Andean slopes, and the Pacific Forest—and likely tied to anthropogenic climate change.

Brower and Garzón-Orduña also explore the history of another widespread group of animals—Neotropical butterflies—but through the use of molecular data. Because the fossil record is too scarce to study the age of origin of these species, the authors estimate time of speciation through phylogenetic trees calibrated with relaxed molecular clocks. Brower and Garzón-Orduña compile clock-based age estimates for six butterfly families, at a Neotropical level, containing ~800 species in ~100 genera. The authors conclude that crown ages, that is, the ages of the initial diversification of the genera, range from Oligocene to Miocene, whereas species pairs diverged mostly during the Pleistocene. The analysis suggest that Neotropical butterflies are relatively young, and that Pleistocene climate changes likely played a significant role, whereas biogeographically significant geological events in the Miocene only set the stage for their subsequent evolution.

While studies on the diversity and community composition of macroorganisms are abundant, this is not the case for microorganisms, especially in the tropics, where this type of surveys are particularly scarce. Benito and Fritz fill this gap in our book, reviewing the diversity and distribution of Neotropical diatoms using a newly created database for tropical South America, which also includes geoclimatic and limnological parameters, as environmental drivers of community change. Using island biogeography and metacommunity approaches, the authors identify several aspects that may influence biogeographical patterns in diatoms, notably differences in regional floras, latitudinal gradients in species richness, dispersal ability, and environmental filtering. They also emphasize the need for incorporating diatoms

into macroecological frameworks for studying community assembly in the Neotropics.

Part II: Regional Biodiversity Patterns and Diversification Processes

This part has 11 chapters that are sorted geographically.

Cracraft et al. revisit Amazonian diversification in relation to the origin of modern landscape features modeled by tectonic activity, drainage evolution, and climatic change. These authors observe that, although current Amazon landscapes were shaped during the last 4–5 million years, much of bird, mammal and butterfly diversity, at species level, originated during the last 1–2 million years. Therefore, the more significant landscape changes for the evolution of the current Amazon biota should have occurred over the last 1–2 million years. Cracraft et al. conclude that many mysteries still remain in relation to the origin of current Amazonian biodiversity, and that further research should be transdisciplinary and integrative, thus avoiding simplistic explanations. Phylogeographic analyses of individual clades should be replaced by among-clade studies that consider a wider spatial context and the differential action of landscape changes on different clades.

To address this question at a smaller geographic scale, Capurucho and collaborators focus on an under-explored region of Amazonia: the White Sand Ecosystems (WSE). These are nutrient-poor, quartz-rich sandy soils patchily distributed in areas of low elevation areas. Gathering data on community composition and the phylogeny of birds and plants, the authors demonstrate that the WSE present a unique assemblage of species. Although these areas pre-date the Quaternary, the molecular data suggest that their Pleistocene history has been dynamic. In the face of anthropogenic change, the authors emphasize that the WSE's low resilience to disturbance, as inferred through their historical analysis, ought to be incorporated into conservation planning.

Many traditional studies of the origin of Amazonian biodiversity, like Cracraft et al. and Capurucho and collaborators', have focused on the action of natural drivers such as tectonic or climatic changes—but largely disregarded the contribution of human activities on local biodiversity patterns. Montoya et al. analyze the potential impact of pre-Columbian anthropogenic activity on today's diversity of Amazonian plants, based on a review of the recent literature. Pre-Columbian human influence began with plant domestication and possibly megafaunal extinction, and remained until European contact, as shown by several types of evidence, including landscape transformations (i.e., savanna expansions) and changes in forest composition and soil properties. However, most studies developed to date are of local nature and are unable to account for the high spatial heterogeneity of the Amazon region. The authors conclude that more transdisciplinary interaction is needed for a proper understanding of human contribution to the Amazonian biodiversity, with emphasis on the last few millennia.

Azevedo and collaborators focus on the history of another key Neotropical ecosystem: the South American savannas. By combining data from the fossil record, paleoenvironmental proxies, and phylogenetic information of plants and animals, the authors claim that the phenomenal diversity of the South American Neotropical

savannas did not emerge from a long history of stability. Paleoenvironmental data indicate that appropriate environmental conditions for the occurrence of savannas were in place since the middle Miocene, in agreement with phylogenetic analyses that recovered that as the beginning of the diversification phase of C4 grasses. However, the review finds that woody plants diversified only a few millions year later. This time lag, say the authors, may indicate that trees adapted to the fire-prone savannas reached ecological dominance only after the emergence of open habitats and their associated fauna.

Colevatti et al. review the biogeographic patterns and the diversification timing of Angiosperms of the South American Dry Diagonal (SADD), an open vegetation belt extending from northeastern Brazil to northern Argentina that includes several xeric biomes. The main goal is to unravel whether the Neogene tectonic/paleogeographic events or the Quaternary climatic changes were the main responsible for biotic diversification. Despite the low proportion of species studied, the authors highlight some regular patterns, notably that major lineage divergences occurred during the Pliocene but most intraspecific divergence took place during the Early and Middle Pleistocene. This suggests that Quaternary climate changes played a significant role but the authors emphasize that species responded differently to these environmental shifts and challenge the hypothesis of glacial refugia and the Last Glacial Maximum (LGM) on the diversification of the SADD Angiosperms.

Guedes et al. discuss the role of the major Non-Andean South American mountain ranges as biodiversity cradles and museums, emphasizing the Serra do Mar Range, the Mantiqueira Mountains, the Espinhaço Mountains, the Northeastern Highlands, the Central Brazilian Highlands, and the Pantepui region. Data published to date show that the Neogene and the Quaternary were important periods for the diversification for many terrestrial groups, resulting in the high endemism that today characterizes the non-Andean mountains. They also discuss how biotic interchange among the mountain ranges themselves, and the Andes, resulted in the patterns observed today.

Peres et al. zooms into the patterns of diversity and endemism of the ecosystem that encompass three of the mountain ranges discussed by Guedes and collaborators: the Brazilian Atlantic rainforest. Their review indicates that higher species richness is often found in the topographically complex coast of Rio de Janeiro and São Paulo, including the Mar and Mantiqueira ranges discussed by Guedes, though not exclusively (e.g. coastal Bahia). Data from multiple groups of plants and animals demonstrate striking differences in the composition of communities in the northern vs. the southern half of the forest, although the exact point of community turnover varies among groups. Several areas of unique taxonomic composition and endemism have been consistently identified across plants and animals, and those defined by organisms with low dispersal ability, such as terrestrial invertebrates, are nested within the areas identified by groups of higher ability to disperse, like birds. By contrasting regional diversity patterns and the processes that led to them, the authors argue that both climate and landscape changed acted in combination to lead to the astonishing diversity of this hotspot. They also argue that while the spatial

patterns of species and genetic diversity are similar across plants and animals, the processes that led to them—and their timing—are widely different.

Silveira et al. argue that the study of Neotropical diversity has largely focused on relatively recent geological environments such as Amazonia or the Andes, whereas older and extremely nutrient-poor settings have been traditionally overlooked, despite their outstanding biodiversity and endemism patterns. This is the case of the so called Old Climatically-Buffered Infertile Landscapes (OCBILs) and, specifically, of the Brazilian “campos rupestres”, which Silveira et al. use to propose a theoretical diversification framework including mechanistic explanations that link ecological and evolutionary processes to vegetation patterns and functional traits. This example is proposed as a model to demonstrate how the OCBIL theory and its predictions can help understanding the complex history of Neotropical plant diversification and inform their conservation.

Rull and Vegas-Vilarrúbia focus on another area included within the OCBIL category, as is the biogeographical province of Pantepui, formed by the assemblage of flat summits of the Guiana table mountains (tepuis), situated in the confluence of the Amazon and the Orinoco basins, between approximately 1500 and 3000 m elevation. Pantepui is one of the few pristine areas remaining on Earth and has been considered a natural laboratory for the study of Neotropical biodiversity. The diversity and endemism patterns of Pantepui are outstanding and its biota and ecosystems are unparalleled across the planet. The authors use the most recent update on the biodiversity of Pantepui, including algae, bryophytes, vascular plants, aquatic insects, butterflies, scorpions, land snails, amphibians, reptiles, birds, mammals and vertebrate parasites, to take the first steps towards a biogeographical, ecological and evolutionary synthesis on the origin of Pantepui biota and ecosystems. Direct (notably fire) and indirect (global warming) threats to Pantepui biodiversity are also analyzed.

Reginato and Michelangeli explore the data emerging from rich, well-curated, and digitized plant collections from Eastern Brazil to advance on the review provided by Peres and collaborators and identify areas of unique community composition (or bioregions). To enable analyses across groups with striking different dispersal abilities, diversity and endemism, they provide three levels of grouping—and the small, intermediate, and large spatial scales. The intermediate-level analysis splits E Brazil into 10 bioregions, which cluster into 5 super-bioregions at the larger scale, and which can be further subdivided into 23 sub-regions, at the smaller scale. High diversity and endemism is flagged both in the mountain ranges of the Serra do Mar and Mantiqueira discussed by Guedes and collaborators, but also in the campos rupestres system introduced by Silveira et al., and lowland areas of Bahia, as described by Peres et al. The rasters of the inferred bioregions are available for download and use by the scientific community at large.

Figueroa-Rangel and collaborators transport us north to Central America. Two major types of Mexican forests have been recognized: the Neartic forests, in the north, and the Neotropical forests, in the south, separated by a transition zone called the Trans-Mexican Volcanic Belt (TMVB). Figueroa-Rangel et al. review the long-term plant diversity changes along an altitudinal transect of the TMVB and the

Neotropical forests, ranging from sea level to the highlands (~3900 m). For that, they use a variety of diversity indices applied to Holocene pollen records. The results reveal that Neotropical forests are more diverse than forests with Holarctic affinities, and that Holocene climate change has been decisive for the shaping of these diversity patterns.

Finally, Roncal et al. discuss an important but often overlooked Neotropical biodiversity hotspot: the Caribbean archipelago. The origins of the high diversity observed in this area is very poorly understood. Roncal et al. use a combination of time-calibrated phylogenetic studies and biogeographical inferences to test the main hypotheses erected to explain Caribbean plant diversity, namely Late Cretaceous vicariance, Oligocene GAARlandia colonization, recurrent Cenozoic long-distance colonization, and in situ speciation. The authors conclude that plants repeatedly colonized the Caribbean islands during the last 60 million years (roughly, the Cenozoic) by long-distance dispersal from the American mainland, especially from Central and South America. They also raise doubts on the existence of GAARlandia, a hypothetical emerged land connecting the present Greater Antilles (Cuba, Hispaniola, Puerto Rico, Jamaica), which would have promoted plant migration among them.

Part III: Taxon-Based Studies at the Local and Regional Scales

This part has eight chapters, each devoted to a unique taxonomic group and its diversification history in a key Neotropical region.

Señaris and Rojas-Runjaic describe and discuss diversity patterns of the herpetofauna (amphibians and reptiles) of the Venezuelan Guayana (also Guiana), one of the largest wilderness areas on the planet. They document how species richness decreases, and endemism increases, with elevation: 34% (amphibians) and 22% (reptiles) of the local fauna is endemic to upland areas, whereas widespread species, or forms with an Amazonian-Guianan distribution, dominate the lowland fauna. This distribution pattern makes conservation particularly difficult in the region, especially in the face of climate change—a topic discussed by the authors.

Exploring a similar topic, Cadena and Céspedes zoom into the processes leading to species turnover along elevational gradients, but they use the Andes as a model system. Combining phylogenetic and occurrence data from a group of passerine birds, they find that most species have small elevational ranges and that those taxa that are more closely related to each other have similar elevational distributions across mountains. Conversely, those species that replace each other along a mountain are distantly related. Based on this pattern, the authors propose that speciation happened mostly in allopatry, across mountains, instead of not along environmental gradients within mountains—suggesting that species accumulation in montane environments largely reflects colonization instead of in situ divergence.

Leubert and collaborators also study patterns of species distributions, this time with the aim of understanding niche evolution in South America. Specifically, they use data from ten plant clades to study a well-documented disjunction: that between the southern Andes (SA) and southern Brazil (SB). These areas tend to have higher precipitation relative to their intervening areas, where no species of the disjunct

clades occur. Comparing divergence times between SA and SB, obtained from phylogenetic studies, they show a wide range of split times across groups—from 27 to 1 Ma. This, they argue, demonstrates that SA-SB disjunctions reflect different clade histories as opposed to a single explanation behind the common pattern observed—much like Peres et al.'s findings for coastal Brazil. Comparing the realized climatic niches of species across SA and SB, however, they find that they do not overlap—suggesting that niche evolution may have occurred across multiple groups. This hypothesis, however, remains to be tested.

Moraes and collaborators transport us to a very distinct environment to investigate the potential role of another environmental barrier on the distribution of local diversity—the Tapajós River, in Amazonia. Using molecular data from 16 species of amphibians and reptiles, sampled from across the Tapajós river basin, the authors find that most groups show spatial patterns of genetic diversity that are consistent with the present-day location of the river: they observe high levels of genetic divergence across river banks. However, much like Leubert et al., the range of riverine-associated divergence times is wide and continuous—suggesting a dynamic history and multiple events of vicariance or colonization.

Recoder and Rodrigues focus on another Neotropical river system to test how changes in geomorphology and climate may have impacted local patterns of diversification—the São Francisco. Located within the domain of the semi-arid Caatinga, the São Francisco river is known for its extensive paleodunes and for the spectacularly diverse sand-dwelling fauna endemic to the dune region. The authors compile existing and new data from local reptiles to ask whether a paleolacustrine vicariant diversification hypothesis, once proposed to explain the origin of this endemic fauna, agrees with current patterns of diversification. That hypothesis suggested that the river flowed into an interior lake during the last glacial maximum, but later made its way out into the Atlantic, isolating populations on opposite margins. The phylogenetic data gathered here show that this model is not sufficient to explain all pattern observed: divergence times differ across species, and some splits pre-date the Pleistocene. Again, Neotropical history is complex and cannot be attributed to a single event.

Ignacio de La Riva calls attention to the diversification processes impacting high-Andean direct-developing frogs, particularly focusing on species occurring from Colombia to Bolivia. Through his review, he clarifies how similar life histories and ecological niches are associated with a particular morphology: local frogs are small-bodied, have short legs, and simple digital tips, likely reflecting adaptations to live in humid mossed habitats of the cloud forests and paramos. According to the author, new studies indicate that this is a convergent morphology across distantly related groups, and that the limited information available about these species render inferences about their evolutionary history, and the history of these unique Neotropical high-elevation areas, tentative at best.

Olave and collaborators also discuss morphological evolution, but are able to incorporate extensive molecular data available for the Liolaemini—the most diverse lizard group of the southern half of South America—in their chapter. Combining molecular and morphological data, they identify different patterns of diversification

among the three genera that compose the clade. They propose that the richness of one of the genera, the species-rich *Liolaemus*, is associated with low extinction rates over time—while the opposite is observed in *Phymaturus*. Based on their analyses, they also argue that natural selection is tied to body size evolution and net diversification in this group, but notice that the trajectories and processes are different across the genera studied.

In the last chapter of Part III, Vargas and Dick focus on a group of 10 genera (ca. 232 species) of Lechytidaceae—the third most abundant family of trees in Amazon forests. Using a time-calibrated phylogeny of more than 100 species, they discuss the diversification history of this important group. Their results suggest that these plants dispersed from the Paleotropics long after the Gondwana break-up, and that most clades diversified in the Miocene. The analysis places the Guayana floristic region as ancestral to several clades, agreeing with Señaris and Rojas-Runjaic about the evolutionary relevance of that area. The results also suggest that this clade diversified through a pattern of steady accumulation of species, as opposed to well-marked pulses in response to specific geomorphological or climatic events.

Epilogue

Closing our book, Carnaval compiles several conservation-relevant messages discussed throughout this volume. Although not the main goal of this contribution, the author argues that one can and should leverage the large, heterogeneous and collective body of knowledge presented in these chapters to inform and guide conservation in the tropical Americas. She emphasizes some of the messages that we are about to hear in the chapters to come, particularly those about the status and patterns of diversity in the Neotropics, along with the authors' insights about the ecological and evolutionary process, and the biological mechanisms, that underscore them. This understanding, she argues, may allow us to better anticipate the behavior of the Neotropical biota as we move forward—a step which will require us not only to integrate data and insight across the biological and physical sciences, as we have done here, but also to reach out to the social sciences. But that is a topic for another book.

References

- Brown JH (2014) Why are there so many species in the tropics? *J Biogeogr* 41:8–22
- Carnaval AC, Waltari E, Rodrigues MT et al (2014) Prediction of phylogeographic endemism in an environmental complex biome. *Proc R Soc B* 281:20141461
- Rull V (2011) Neotropical biodiversity: timing and potential drivers. *Trends Ecol Evol* 26:508–513